## **Workers Comp**

Date Ph: (214) 919-2090 or (877) 753-6878 Fax: 1 (888) 294-9434 ☐ New Patient DOB Patient Name Street Address: Apt # \_\_\_\_City\_\_\_\_ Zip State Phone #\_\_\_\_\_ Cell # Allergies ICD 9 Code: □ 4 mg □ 8 mg □ Zofran ☐ Ibuprofen □ 800mg □ 600mg □ Phenergan □ 25 mg □ 50 mg □ Naproxen ☐ 375mg □ 500mg ☐ Etodolac □ 300mg ☐ 400mg □ 500mg Qty:\_ Refills ☐ Meloxicam ☐ 7.5mg □ 15mg` □ Amitriptyline □ 10mg □ 25mg □ 50mg □ 75mg Sig: Refills Qty: ☐ 100mg ☐ Cymbalta □ 20mg □ 30mg □ 60mg ☐ Venlafexine ER ☐ 37.5mg ☐ 75mg □ 150mg ☐ Gabapentin ☐ 100mg ☐ 300mg ☐ 400mg, Refills Sig:\_ Qty: □ 600mg □ 800mg □ 10mg ☐ Cyclobenzaprine ☐ 5mg Qty: Refills Sig:\_\_\_\_ ☐ 2mg ☐ Tizanidine ☐ 6mg 4mg ☐ Methylprednisone Dosepack ☐ 4mg ☐ Prednisone ☐ 5mg ☐ 10mg ☐ 20mg □ 750mg ☐ Methocarbamol □ 500mg Sig: Qty:\_ Refills Refills Sig: Qty: □ 100mg ☐ Celebrex □ 200mg ☐ 400mg ☐ **Famotidine** ☐ 20mg ☐ 40mg Qty:\_\_\_\_\_ Refills Refills Sig: Sig: Qty: R ALL controlled substance quantities must be hand written in number and letter form ADDITIONAL INJURED WORKER MEDICATION ☐ Capsaicin 0.038% + Menthol 5% (For Pain) **OSTEOARTHRITIS:** ☐ Lidocaine HCL 4% + Menthol 1% (For Pain) \*Dosing = Take 1 capsule 3 times daily Synovacin \*Dosing = Apply 1 - 3 patches to affected area daily ☐ Genicin QTY:  $\square$  30 (Thirty)  $\square$  60 (Sixty)  $\square$  90 (Ninty) QTY: 90 (Ninty) 180 (One-Hundred Eighty)
Refills: 1 (One) 2 (Two) 3 (Three) Refills: 1 (One) ☐ 2 (Two) ☐ 3 (Three) PAIN AND INFLAMATION: STOOL SOFTNER: ☐ Theramine ☐ 120 \*Dosing = Take with full glass of water (8oz). For dietary management of pain syndromes that include chronic pain, fibromyalgia, neuropathic pain, and inflammatory pain Do not take medicine more than directed. Take two (2) capsules twice daily or as needed under physicain supervision ☐ Laxacin ☐ Trepadone ☐ 120 ☐ **Docusate 100mg** Up to QID QTY: 100 Refills: 1 (One) 2 (Two) 3 (Three) For the dietary management of pain associated with osteoarthritis Take two (2) capsules twice daily SLEEP DISORDER: ☐ Somnicin Refills:  $\square$  1 (One)  $\square$  2 (Two)  $\square$  3 (Three) \*Dosing = Take 1 tablet 30 minutes prior to bedtime ☐ GABAdone ☐ 60 ☐ SentraPM For the dietary management of sleep disorders associated with anxiety \*Dosing = Take 1 or 2 capsules 30 minutes prior to bedtime Take one (1) or two (2) capsules at bedtime to initiate sleep QTY:  $\square$  30 (Thirty)  $\square$  60 (Sixty)  $\square$  90 (Ninty) QTY:\_\_\_\_\_ Refills: \_\_\_\_\_  $\square$  1 (One)  $\square$  2 (Two)  $\square$  3 (Three) Refills: ☐ 1 (One) ☐ 2 (Two) ☐ 3 (Three) 
 Prescriber's Name
 Office Contact

 Street Address
 Suite #
 City
 State
 Zip
 License # NPI # DPS # DEA # Telephone

IMPORTANCE NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged, proprietary or exempt from disclosure under applicable laws. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately. Medicare and Medicaid or another state funded program will not cover above mentioned products. Co-payments due at dispensing of the medication

**Prescriber's Signature** (signature required. NO STAMPS)

FDA BLACK BOX WARNING: [NSAID] may cause an increased risk of serious cardiovascular thrombotic event, myocardialinfarction, and stroke, which can be fatal. This risk may increase with duration of use. Patients with cardiovascular diesase or risk factors for cardiovascular diesase may be at greater risk.